

University of Groningen

Highly enantioselective DNA-based catalysis

Roelfes, Gerard; Boersma, Arnold J.; Feringa, Bernard

Published in:
ChemInform

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2006

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Roelfes, G., Boersma, A. J., & Feringa, B. (2006). Highly enantioselective DNA-based catalysis. ChemInform, 37(25).

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Supplementary Information Accompanying

“Highly Enantioselective DNA-based Catalysis”

by

Gerard Roelfes, Arnold J. Boersma and Ben L. Feringa

Figure S1. Induced CD spectra from $[\text{Cu}(\text{L}_{4-7})(\text{NO}_3)_2]$ (150 μM) combined with salmon testes DNA (0.8 mg/ml) in Mops buffer (20 mM, pH 6.5). a) only st-DNA; b) $[\text{Cu}(\text{bipy})(\text{NO}_3)_2]$ / st-DNA; c) $[\text{Cu}(\text{dppz})(\text{NO}_3)_2]$ / st-DNA; d) $[\text{Cu}(\text{dpq})(\text{NO}_3)_2]$ / st-DNA; e) $[\text{Cu}(\text{phen})(\text{NO}_3)_2]$ / st-DNA.




Table S1: Effect of variation of the concentration of $[\text{Cu}(\text{bipy})(\text{NO}_3)_2]$, st-DNA and **2a** on the results of the catalyzed reaction.

Entry	Concentration $[\text{Cu}(\text{bipy})(\text{NO}_3)_2]$ (mM)	Conc. DNA (mg/ml)	dienophile (mM)	Conversion (%)	Endo:Exo	E.e. Endo
1	0.05	1.3	1	>80	97:3	89
2	0.10	1.3	1	>80	98:2	91
3	0.15	1.3	1	>80	98:2	91
4	0.30	1.3	1	>80	98:2	89
5	0.45	1.3	1	>80	98:2	85
6	0.60	1.3	1	>80	98:2	77
7	0.15	0.65	1	68	98:2	89
8	0.30	0.65	1	>80	98:2	84
9	0.30	1.3	0.3	Quant.	97:3	90
10	0.30	1.3	1.5	>80	98:2	89
11	0.30	1.3	3	>80	97:3	87
12	0.30	1.3	6	65	97:3	86

Table S2

	catalyst	T	endo:exo	e.e. endo (%)
1^a	Cu(NO ₃) ₂ / DNA	RT	95:5	10
2^b	DNA	5 °C	n.d. ^c	<5 %
3^d	[Cu(dppz)(NO ₃) ₂] / DNA	5 °C	94:6	-

Table S2, Selection of relevant control experiments, performed under standard conditions. a) conversion 50-60 %. b) conversion < 5%. c) cannot be determined due to the low conversion. d) conversion 52 %.

Experimental and Synthetic Procedures

General remarks

Salmon testes and calf thymus DNA were obtained from Sigma.

Physical methods.

Equilibrium binding constants to salmon testes DNA were determined by UV/Vis titration, following the procedure of Meehan.¹ After dissolution of salmon testes DNA (2 mg/ml), the stock solution was dialyzed extensively against Mops buffer (20 mM pH 6.5) prior to use. The concentration in base pairs was determined spectrophotometrically, using $\epsilon_{260} = 12800 \text{ M}^{-1} \text{ cm}^{-1}$.² The absorbance ratio of $\lambda_{260}/\lambda_{280}$ was 1.8-1.9, indicating the DNA was sufficiently free of protein. The K_b was determined by titration of DNA to a solution of copper complex in buffered solution. Concentrations of copper complexes generally were 30 μM , or 15 μM in case of Cu(dppz)(NO₃)₂ and Cu(dpq)(NO₃)₂. Under conditions where the ratio of bound complex : DNA base pairs approaches zero, the K_b can be determined using :

$$\frac{D}{\Delta\epsilon_{ap}} = \frac{1}{\Delta\epsilon} D + \frac{1}{\Delta\epsilon K_b}$$

where $\Delta\epsilon_{ap} = |\epsilon_a - \epsilon_f|$, $\Delta\epsilon = |\epsilon_b - \epsilon_f|$, ϵ_a , ϵ_f and ϵ_b are the apparent, free and bound extinction coefficients for the complex, respectively, and D is the DNA concentration in basepairs. In a plot of $D/\Delta\epsilon_{ap}$ vs. D , K_b is given by the ratio of the slope to the y intercept. A representative plot, for [Cu(bipy)(NO₃)₂], is shown below.




Figure S2. Representative plot of $[DNA]/\Delta\epsilon_{ap}$ vs. $[DNA]$ for [Cu(bipy)(NO₃)₂] (30 μM). The solid line represents the least squares linear fit of the data.

Catalytic Diels-Alder reactions, representative procedure.

A buffered solution (20 mM Mops, pH 6.5) of DNA bound catalyst (1.3 mg/ml salmon testes DNA and 0.3 mM $[\text{Cu}(\text{bipy})(\text{NO}_3)_2]$) was prepared by mixing a solution of salmon testes DNA (10 ml of a 2 mg/ml solution in 30 mM Mops, prepared 24 h in advance) with an aqueous solution of catalyst (5 ml of a 0.9 mM solution, prepared by adding a solution of $[\text{Cu}(\text{bipy})(\text{NO}_3)_2]$ in a minimal amount of dmso to 5 ml H_2O). An aliquot of a stock solution of dienophile **2a** in CH_3CN (30 μL of a 0.5 M soln., final conc. 1 mM) was added and the mixture was cooled to 5 $^\circ\text{C}$. The reaction was started by addition of cyclopentadiene (21 μL , final conc. 15 mM) and mixed by continuous inversion for 3 days, followed by extraction of the product with diethyl ether. After H-NMR analysis the e.e. was determined by chiral hplc (Daicel chiralcel-ODH column). Selected products were purified by column chromatography and analyzed on a Daicel chiralcel-ODH column or Daicel chiralpak-AD column to confirm the results obtained from analysis of the crude product.

HPLC conditions:

3a: Daicel chiralcel-ODH, heptane/*i*PrOH 98:2, 0.5 ml/min. Retention times: 11.4, 12.3 (exo isomer), 13.7 and 16.7 mins (endo isomer); Daicel chiralpak-AD, heptane/*i*PrOH 99:1, 1 ml/min. Retention times: 9.7, 10.6 (exo isomer), 12.5 and 14.6 mins (endo isomer).

3b: Daicel chiralpak-AD, heptane/*i*PrOH 98:2, 1 ml/min. Retention times: 13.5, 15.2 (exo isomer), 17.4 and 21.1 mins (endo isomer).

3c: Daicel chiralcel-ODH, heptane/*i*PrOH 99.75:0.25, 0.5 ml/min. Retention times: 13.9, 14.7 (exo isomer), 20.4 and 25.3 mins (endo isomer).




Figure S3. ^1H -NMR of the crude product **3a** of the Diels-Alder reaction catalyzed by $[\text{Cu}(\text{dmbipy})(\text{NO}_3)_2]$ (table 1, entry 16). S denotes residual starting material **2a**.




Figure S4. HPLC trace of the crude product **3a** of the Diels-Alder reaction catalyzed by $[\text{Cu}(\text{dmbipy})(\text{NO}_3)_2]$ (table 1, entry 16). S denotes residual starting material **2a**. The peak of one of the enantiomers of the exo product (at 11.4 min) is too small to detect.

Synthesis

General remarks

Dienophiles **2a-b**,³ $\text{Cu}(\text{dppz})(\text{NO}_3)_2$,⁴ $\text{Cu}(\text{dpq})(\text{NO}_3)_2$,⁴ 2-(2-pyridyl)imidazole (**9**)⁵ were prepared following published procedures.

(E)-4,4-dimethyl-1-(2-pyridinyl)-2-penten-1-one (2c). This compound was prepared following the procedure as described for **2a**.³ Starting from 2-acetylpyridine (2.06 g, 17 mmol) and pivaldehyde (1.42 g, 16.5 mmol), after column chromatography (SiO_2 , heptane/ethyl acetate 8:1), **2c** was obtained as a white solid. Yield: 706 mg, 3.7 mmol, 22 %. $^1\text{H-NMR}$ (CDCl_3 , 400 MHz) δ 1.18 (s, 9H), 7.24 (dd, 1H, $J = 15.8$ Hz, $J = 0.8$ Hz), 7.46 (m, 1H), 7.54 (d, 1H, $J = 16.5$ Hz), 7.84 (m, 1H), 8.12 (d, 1H, $J = 7.3$ Hz), 8.71 (m, 1H). $^1\text{H-NMR}$ (CDCl_3 , 100 MHz) δ 28.71 (q), 34.39 (s), 119.32 (d), 122.81 (d), 126.65 (d), 136.85 (d), 148.76 (d), 154.29 (s), 159.89 (d), 190.04 (d); Anal. Calcd for $\text{C}_{12}\text{H}_{15}\text{NO}$: C, 76.16 H, 7.99 N, 7.40. Found: C, 76.1 H, 8.04 N, 7.45.

[3-(tert-butyl)bicyclo[2.2.1]hept-5-en-2-yl](2-pyridinyl)methanone (3c, major isomer)

$^1\text{H-NMR}$ (CDCl_3 , 400 MHz) δ 0.89 (s, 9H), 1.38 (m, 1H), 1.80 (m, 1H), 1.90 (dd, 1H, $J = 6.2$ Hz, $J = 1.5$ Hz), 2.76 (dd, 1H, $J = 2.9$ Hz, $J = 1.5$ Hz), 3.22 (m, 1H), 4.26 (dd, 1H, $J = 6.2$ Hz, $J = 2.9$ Hz), 5.69 (dd, 1H, $J = 5.5$ Hz, $J = 2.6$ Hz), 6.43 (dd, 1H, $J = 5.5$ Hz, $J = 2.9$ Hz), 7.44 (m, 1H), 7.78 (m, 1H), 7.97 (d, 1H, $J = 7.7$ Hz), 8.71 Hz (d, 1H, $J = 4.8$ Hz); MS (CI): 256 (M+1); HRMS Calcd for $\text{C}_{17}\text{H}_{21}\text{N}_1\text{O}_1$ 255.1623, found 255.1613.

$[\text{Cu}(\text{phen})(\text{NO}_3)_2]$. Following the procedure as described for $[\text{Cu}(\text{dppz})(\text{NO}_3)_2]$,⁴ starting from phenanthroline (70 mg, 0.35 mmol) and $\text{Cu}(\text{NO}_3)_2 \cdot 3\text{H}_2\text{O}$ (94 mg, 1.1 eq), $[\text{Cu}(\text{phen})(\text{NO}_3)_2]$ was obtained as a blue solid. Yield: 114 mg, 0.31 mmol, 89 %. Anal. Calcd for $\text{C}_{12}\text{H}_8\text{CuN}_4\text{O}_6$: C, 39.19 H, 2.19 N, 15.23. Found: C, 39.25 H, 2.09 N, 15.15.

$[\text{Cu}(\text{bipy})(\text{NO}_3)_2]$. Following the procedure as described for $[\text{Cu}(\text{dppz})(\text{NO}_3)_2]$, starting from 2,2'-bipyridine (60 mg, 0.39 mmol) and $\text{Cu}(\text{NO}_3)_2 \cdot 3\text{H}_2\text{O}$ (100 mg, 1.1 eq),⁴ $[\text{Cu}(\text{bipy})(\text{NO}_3)_2]$ was obtained as a blue solid. Yield: 86 mg, 0.25 mmol, 64 %. Anal. Calcd for $\text{C}_{10}\text{H}_8\text{CuN}_4\text{O}_6$: C, 34.94 H, 2.35 N, 16.30. Found: C, 35.1 H, 2.30 N, 16.15.

$[\text{Cu}(\text{2-(2-pyridyl)imidazole})(\text{NO}_3)_2 \cdot \text{H}_2\text{O}]$. To a solution of 2-(2-pyridyl)imidazole (**9**) (74 mg, 0.51 mmol) in ethanol (10 mL) was added $\text{Cu}(\text{NO}_3)_2 \cdot 3\text{H}_2\text{O}$ (123 mg, 0.51 mmol). The mixture was shaken until a clear green solution was obtained. The solution was placed in an ether bath for 2 days. The green crystals were filtered and washed with water and ethanol. Yield: 65 mg, 38%. Anal. Calcd for $\text{C}_8\text{H}_9\text{CuN}_5\text{O}_7$: C, 27.40 H, 2.59 N, 19.97. Found: C, 27.6 H, 2.49 N, 19.74.

[Cu(2-(2-pyridyl)benzimidazole)(NO₃)₂·H₂O]. To a solution of Cu(NO₃)₂·3H₂O (97 mg, 0.39 mmol) in a mixture of acetone (4 mL) and ethanol (0.3 ml) was added a solution of 2-(2-pyridyl)benzimidazole (**10**) (75 mg, 0.38 mmol) in ethyl acetate (4ml), through a small cotton plug. The dark green solution was filtered and the vial was closed with a cotton plug, allowing for slow evaporation of acetone. After one night a dark green solid had precipitated, which was washed with a small volume of ethyl acetate. Yield: 135 mg, 89 %. Anal. Calcd for C₁₂H₁₁CuN₅O₇: C, 35.96 H, 2.77 N, 17.47. Found: C, 36.30 H, 2.83 N, 17.04.

[Cu(4,4'-dimethyl-2,2'-dipyridyl)(NO₃)₂]. To a solution of Cu(NO₃)₂·3H₂O (0.10 g, 0.41 mmol) in ethanol was added 4,4'-dimethyl-2,2'-dipyridyl (**11**) (38 mg, 0.24 mmol), dissolved in ethanol. The solution was placed in an ethyl acetate bath and left standing for 2 days. The blue solid was filtered and washed with ethanol. Yield: 49 mg, 59%. Anal. Calcd for C₁₀H₈CuN₄O₆: C, 38.8 H, 3.25 N, 15.07. Found: C, 38.5 H, 3.14 N, 14.80.

Reference List

1. A. Wolfe, G. H. Shimer, T. Meehan, *Biochemistry*, **1987**, 26, 6392-6396.
2. F. Barcelo, D. Capo, J. Portugal, *Nucl. Acids. Res.*, **2002**, 30, 4567-4573.
3. S. Otto, J. B. F. N. Engberts, *J. Am. Chem. Soc.*, **1999**, 121, 6798-6806.
4. M. Navarro, E. J. Cisneros-Fajardo, A. Sierralta, M. Fernandez-Mestre, P. Silva, D. Arrieche, E. Marchan, *J. Biol. Inorg. Chem.*, **2003**, 8, 401-408.
5. K. A. Reeder, E. V. Dose, L. J. Wilson, *Inorg. Chem.*, **1978**, 17, 1071-1075.